

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/40, 9/14, 9/48, 9/52, 9/62		A1	(11) International Publication Number: WO 98/50019 (43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/09449 (22) International Filing Date: 8 May 1998 (08.05.98) (30) Priority Data: 60/046,089 9 May 1997 (09.05.97) US 08/950,432 15 October 1997 (15.10.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 60/046,089 (CIP) Filed on 9 May 1997 (09.05.97) US 08/950,432 (CIP) Filed on 15 October 1997 (15.10.97) (71) Applicant (for all designated States except US): SAGE PHARMACEUTICALS, INC. [US/US]; 5408 Interstate Avenue, Shreveport, LA 71109 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): CHEN, Jivn-Ren [US/US]; 7614 Brook Haven Way, Shreveport, LA 71105 (US). (74) Agent: CHAN, Albert, Wai-Kit; Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, NY 10036 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: STABLE ORAL PHARMACEUTICAL DOSAGE FORMS			
(57) Abstract The present invention relates to new stable enteric coated pharmaceutical dosage forms for oral use containing Omeprazole or Lansoprazole, to a formulation and a method for the manufacture of such a dosage form, and to a method of gastric acid pump inhibition and providing gastrointestinal cytoprotective benefit by using them.			

achieve
oil
alkaline ag
surfactant
suspending ag.

6096543
6077541

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

STABLE ORAL PHARMACEUTICAL DOSAGE FORMS

5 This application is a continuation-in-part application of U.S. Serial No. 08/950,432, filed October 15, 1997, claiming priority of U.S. Provisional Application No. 60/046,089, filed May 9, 1997, the content of which is incorporated by reference into this application.

Background of the Invention

10

U.S. patent number 4,255,431 describes a compound 2-[2-(3,5-dimethyl -4-methoxy)-pyridyl methyl sulfinyl]-(5-methoxy)-benzimidazole (Omeprazole) or pharmaceutically acceptable salt or non-toxic acid addition salt as a therapeutic compound for mammals including man, suffering from gastric acid secretion disturbances.

15

Omeprazole, however is only stable in basic pH conditions and degrades rapidly in neutral or acid pH environment. For this reason the omeprazole oral dosage form must be protected, not only from the acidic inert ingredients used to make a dosage form but also from the acidic gastric fluid in order to intactly reach the absorption site in the small intestine.

20

25

A study [Drug Dev. Ind. Pharm. 21(8), 965 (1995)] showed the effect of pH on the stability of omeprazole solution. The pH-decomposition rate profile curve indicated that the maximum stability was at pH 11. Below pH 7.8 the decomposition was very fast.

30

A survey of the stability of Omeprazole products from 13 countries was reported [Drug Dev. Ind. Pharm. 22(12), 1173(1996)]. The results of this independent survey of the stability of omeprazole solid dosage forms (20 mg) show that product available in many countries worldwide exhibits a very wide range of stability characteristics. Only 18% of

35

-2-

total products tested (34) were considered to demonstrate good physical and chemical stability over the course of the study.

5 U.S. patent number 4,628,098 discloses that:

Lansoprazole is a substituted benzimidazole 2-[[[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridyl] methyl] sulfinyl] benzimidazole, a compound and a pharmacologically acceptable salt thereof that inhibits gastric acid secretion.

10

Lansoprazole is relatively stable when exposed to light. The compound degrades in aqueous solution, the rate of degradation increasing with decreasing pH.

15

It is well known in the pharmaceutical industry that an enteric coating technology is the most efficient means to protect acid unstable medication from the attack of the gastric fluid and can rapidly release the active drug in the proximal part of the gastrointestinal canal.

20

An enteric coated dosage form of omeprazole was reported in 1985 by Pilbrant and Cederberg (Scand. J. Gastroenterology 1985; 20 (supp.108) p. 113 - 120). It was found later that the stability of this dosage form was not sufficient for a long term commercial purpose.

25

Because of the acidic property of the enteric coating polymer, the omeprazole or lansoprazole will degrade and diminish its therapeutic value by direct or indirect contact with it during or after the coating process, even in the presence of alkaline inert ingredients with the active drug in the core particles.

30

DE - AI number 3046 559 teaches a two layer coating system for a dosage form in which the first coating is a water insoluble layer of microcrystalline cellulose and then

35

-3-

enteric coating as the second layer to release the active drug in the colon.

5 WO number 85/03436 describes a technique to mix active drug with buffering components (i.e. sodium dihydrogen phosphate) to maintain a constant pH in a core for particular purposes. The core is coated with a layer which controls the diffusions.

10 All of these inventions when applied to omeprazole's or lansoprazole's case will not give either an adequate release of active drug or storage stability of such dosage form.

15 U.S. patent number 4,786,505 teaches an art to make an oral pharmaceutical preparation comprising (1) a core consisting of omeprazole plus an alkaline reacting compound (2) an inert subcoating of core (3) an enteric coating of subcoated core and to use it in the treatment of
20 gastrointestinal disease.

This technique requires a series of cumbersome and laborious pharmaceutical processes: 1) preparation of the core, which should be suitable for multilayer coating 2)
25 Coating the core with one or more layers of an inert subcoating containing one or more alkaline substances 3) applying an outer enteric coating to the subcoated core 4) make the pharmaceutical preparation for therapeutical use.

30 U.S. patent number 5,232,706 claims an oral pharmaceutical preparation of omeprazole or an alkali salt of omeprazole and a process for producing such preparation. The design principle of the preparation is basically similar to the U.S. patent number 4,786.505. This preparation is
35 comprised of: 1) a nucleus of active drug and first basic organic compound; 2) a first coating of nucleus containing at least a layer of a basic water soluble excipient and a

-4-

second basic organic compound; and 3) a second coating formed by an enteric coating.

5 The major difference between U.S. patent number 5,232,706 and U.S. patent number 4,786,505 is the type of basic compound used; the former uses basic organic compound the later uses inorganic alkaline reaction compounds in core and in subcoating.

10 U.S. patent number 5,385,739 relates to a stable formulation of Omeprazole microgranules containing a neutral core of sugar and starch coated with an active layer of drug and mannitol powder mixture with the aid of a solution of a binding agent in water plus ethanol. An
15 additional protective layer of mannitol and sugar syrup is then applied prior to the final gastroprotection coating.

20 The art of coating a powder mixture containing an active ingredient onto the core of sugar and starch by using a binding solution is a rather difficult process to obtain uniform drug core granules.

25 In addition, two applications of aqueous solution involving the direct contact with Omeprazole, in the coating process, may effect the stability of this moisture-sensitive drug.

30 U.S. patent number 5,399,700 teaches a method for stabilizing an acid unstable benzimidazole derivative, by forming an inclusion complex of Omeprazole with cyclodextrin.

35 This inclusion complex was synthesized in an aqueous alkaline solution at 40° - 70°C and then used to manufacture a tablet which is first coated by a water soluble substance and then with an enteric substance to form an enteric coated oral drug.

-5-

The safety of cyclodextrin for human oral dosage form needs to be clarified. Even if the acid stability and dissolution characterization of this inclusion complex has been well demonstrated in vitro, its behavior in vivo is not referenced in this patent.

U.S. patent number 4,689,333 claims a pharmaceutical composition for preventing or treating digestive ulcers or gastritis which contains an effective amount of Lansoprazole or a pharmacologically acceptable salt thereof and pharmacologically acceptable carriers.

A study [Drug Dev. Ind. pharm. 18 (13), 1437 (1992)] reported that enteric granule formulations for Lansoprazole was established, however, it was difficult because some of the excipients needed for these formulations are incompatible with the drug. The alkaline stabilizers were then screened and the optimal pH stability of 9 was suggested in this paper.

A continued study from the above mentioned effort [Drug Dev. Ind. pharm. 20(9), 1661 (1994)] attempted to prepare more stable enteric granules without using these troublesome excipients by using a centrifugal fluid-bed granulator instead of an extruder-spheronizer. It was said that with this method more stable enteric granules could be obtained. The utilization of special sophisticated equipment to achieve the stable granules may suffer economical disadvantages.

U.S. patent number 5,026,560 discloses spherical granules having a seed core coated with a binder and spraying powder containing lansoprazole as active drug, low substituted hydroxypropylcellulose and magnesium or calcium carbonate as alkaline agents. The powder coated core is further coated with spraying powder of low substituted hydroxypropylcellulose and then with enteric coating agent.

-6-

Technically, powder coating in pharmaceutical manufacturing brings forth issues such as content uniformity of the active on the seed core, laborious process attention and time consumption. It therefore tremendously increases the cost of the product.

U.S. patent number 5,045,321 describes a pharmaceutical composition for coated tablets or granules, which is comprised of lansoprazole being in contact with at least one of the basic inorganic salts evenly. No protective and/or enteric coating is mentioned in the patent claim.

U.S. patent number 5,093,132 is similar to the U.S. patent number 5,045,321 but more specifically, describes an oral stabilized pharmaceutical composition for the inhibition of gastric acid secretion comprising of lansoprazole or its pharmaceutically acceptable salt in contact evenly with a basic inorganic salt stabilizing agent. It also mentions simply an enteric coating for the composition.

Enteric coated oral solid dosage forms have been in existence for a century. U.S. patent number 3,656,997 and 3,959,540 disclosed a coated gastric juice resistant capsules and process for the production thereof. Most commercially available products under this category are coated pellets or granule filled capsules and tablet dosage forms. The concept of an enteric coated capsule dosage for commercial purposes are relatively new. So far, only one prescription product - VIVOTIF BERNA™ (Berna Products, Co., Coral Gables, Fl.) is in the U.S.A. market.

The advantage of enteric coated capsules are duplex. In addition to the merit natures of enteric coating polymers, the problem involved and the need for extensive efforts in development and the preparation of enteric pellets or tablets can be avoided.

-7-

Historically, enteric coated hard shell capsules were manufactured with formaldehyde treatments. This technology results in a stability problem with this product. Since the successful introduction of a number of enteric polymeric materials, the technology of this coating becomes more popular and is increasingly used in the pharmaceutical industry. However, the use of organic solvents causes risks of air pollution and inflammability. With the advent and availability of aqueous enteric polymer dispersions, manufacturing of enteric coated dosage forms have acquired a great deal of attention. The modern coating equipment now offers to pharmaceutical manufacturers better conditions for the application of aqueous coatings.

The process of enteric coating for pharmaceutical dosage forms are not significantly different with other coating processes. Generally, the sample is first placed in the coating pan or in the fluid-bed chamber for fluidization, while a coating solution is then spraying through a gun according to the two operations: 1) spraying or wetting and 2) drying, which is repeated consecutively during the process.

-8-

Summary of the Invention

5 It is well known that the stability of omeprazole and lansoprazole may be affected adversely in the presence of water and organic solvent, especially by the former.

10 The previous art of U.S. patent number 4,786,505 and number 5,232,706 all use conventional aqueous wet technologies to make core pellets or granulations. This invention, by preferably using dry mixing of drugs with alkaline substances as core granulations for either capsule filling or direct tablet compression in its processes, can substantially eliminate the stability risk of the active by excluding the moisture from the core granulation.

15 The alkaline substances in contact with the enteric coating in the aqueous environment can reduce the acid resistance properties of the latter. The invention of the above mentioned two U.S. patents includes the alkaline substances in the subcoating which is directly in contact with the enteric coating, thus it could be detrimental to the gastric fluid resistance of the enteric coating and result in the inferior therapeutic efficacy of omeprazole or lansoprazole.

25 For this reason the present invention uses a capsule shell as a separating barrier between the alkaline core of active drug and the enteric coatings. In the case of a tablet, only the pure, non-ionic polymer is used as a protective coating.

30

35 Furthermore, the present invention describes an improved oral pharmaceutical dosage form for omeprazole, its salt, lansoprazole or its salts. It is comprised of 1) a core powder or granules of active drug and alkaline inorganic or organic substances is formed by dry mixing; 2) said powder or granulation is filled into a hard gelatin capsule or is

-9-

further mixed with other pharmaceutical excipients for direct tablet compression; 3) (for tablet only) a conventional pharmaceutical protective coating is disposed on the tablets; and 4) an outer layer of enteric coating is disposed on the dosage form (capsule or tablet).

This improved dosage form is more economically feasible in terms of time process and material savings. It is sufficiently stable for commercial distribution and storage. It also effectively protects the active drug from the attack of the acidic gastric fluid and efficiently delivers the active drug to the small intestine.

A process for the manufacturing of an oral dosage form of omeprazole or lansoprazole was also described. The details of granulation, encapsulation, tableting and coating art can be found in "The Theory and Practice of Industrial Pharmacy, 1986, Lea & Febiger, Philadelphia, PA, USA, the content of which is incorporated into this application as a reference.

-10-

Detailed Description of the Invention

5 This is an invention of an improved pharmaceutical dosage
form for oral administration to human being or animal host
which is comprised of: (a) a core granulation formed by dry
mixing an acid-unstable drug or its salt and an alkaline
substance or pharmaceutical excipient without using an
aqueous granulating solution; (b) said dry core granulation
10 can then be quantitatively filled into a proper size empty
hard gelatin capsule shell using this shell as a barrier
and hence eliminating the process of applying a protective
coating layer onto the granulation. In other words, this
hard gelatin capsule shell constitutes simultaneously a
barrier between said core granulation and the outer enteric
15 coating of said capsule during the processing to complete
the capsule dosage form; and (c) an enteric coating can
then be applied onto said capsule to prevent the capsule
from releasing the acid-unstable drug in a low pH
environment (i.e. stomach) and then to deliver the drug in
20 a higher pH environment (i.e. small intestine).

As used herein, core granulation is a mixture of a
pharmaceutically acceptable granulated alkaline substance
and, or excipient, and a drug active ingredient that can be
25 processed into uniform spherelike or regularly shaped
aggregates for the improvement of flowability and
compressibility. The manufacturing processes may employ
one, or a combination of, four established methods:

- 30 1) dry mixing;
2) direct compression;
3) milling; and
4) non-aqueous granulation.

35

These methods are described in "The Theory and Practice of
Industrial Pharmacy, 1986, Lea & Febiger, Philadelphia, PA,

-11-

USA, edited by Lachman, L., Lieberman, H.A., and Kanig, J.L.

5 This invention also includes an improved pharmaceutical dosage form for oral administration to human being or animal host which consists of: (a) a core granulation formed by dry mixing an acid-unstable drug or its salt, and an alkaline substance or pharmaceutical excipient, without using an aqueous granulating solution, and which can be
10 directly compressed into a tablet; (b) said tablet can then be filled into an empty, hard gelatin capsule shell using this shell as a barrier and hence eliminating the process of applying a protective layer onto the tablet. In other words, this hard gelatin capsule shell constitutes
15 simultaneously a barrier between the said tablet and an outer enteric coating of said capsule during the processing to complete the capsule dosage form; and (c) an enteric coating can then be applied onto said capsule to prevent the capsule from releasing the acid-unstable drug in the
20 low pH environment (i.e. stomach) and then to deliver the drug in a higher pH environment (i.e. small intestine).

Also included in this invention is an improved pharmaceutical dosage form for oral administration to human
25 being or animal host which consists of: (a) a core granulation formed by dry mixing an acid-unstable drug or its salt, and an alkaline substance or pharmaceutical excipient, without using an aqueous granulating solution, which can then be directly compressed into a tablet; (b)
30 said tablet can then be coated with a non-ionic protective coating in an organic solvent as a barrier: (1) to separate the acid-unstable active drug in the tablet from the outer enteric coating and, (2) to protect the outer enteric coating from the permeation of generated alkaline solution
35 formed by any existed water in the core tablet; and (c) an enteric coating then can be applied onto said tablet to protect it from releasing the acid-unstable drug in a low

-12-

pH environment (i.e. the stomach) and to deliver the active in a higher pH environment (i.e. the small intestine).

5 The protective coating can be applied by a standard film coating procedure in a suitable coating machine using a non-aqueous solution. The non-ionic protective polymer is selected from the group consisting of hydroxypropyl methylcellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, and polyvinylpyrrolidone. The organic solvent
10 is selected from the group consisting of isopropyl alcohol, methanol and ethanol. Other appropriate non-ionic protective polymer may be similarly used.

15 The plasticizer for protective coating includes, but is not limited to, triethyl citrate, propylene glycol and polyethylene glycol 6000.

20 In a preferred embodiment, the drug active ingredient comprises Omeprazole, salt of Omeprazole - selected from the group consisting of sodium, potassium, calcium and ammonium salts, Lansoprazole or salt of Lansoprazole.

25 In another embodiment, the alkaline substance of dosage form mentioned above comprises one or any combination of the following: (a) alkaline metallic salt of carbonic acid: calcium carbonate, granulated calcium carbonate; (b) dicalcium phosphate anhydrous, Dibasic sodium phosphate anhydrous, tricalcium phosphate, anhydrous; (c) sodium carboxymethylcellulose, calcium carboxymethylcellulose; (d)
30 magnesium aluminum silicate; (e) sodium lauryl sulfate; (f) sodium bicarbonate; and (g) microcrystalline cellulose, hydroxypropylmethylcellulose, hydroxyethylcellulose.

35 Also in another embodiment, the pharmaceutical excipient mentioned in the above dosage forms may be selected from one or any combination of the group consisting of dextrose, sorbitol, mannitol, starch, dextrin, maltodextrin, lactose,

-13-

magnesium stearate, calcium stearate, talc, microcrystalline cellulose, hydroxypropylmethylcellulose and hydroxyethylcellulose.

5 Also in another embodiment, the pharmaceutical excipient mentioned in the above dosage forms may be selected from one or any combination of the group consisting of dextrose, sorbitol, mannitol, starch, dextrin, maltodextrin, lactose, magnesium stearate, calcium stearate, talc,
10 microcrystalline cellulose, hydroxypropylmethylcellulose and hydroxyethylcellulose. Other appropriate excipients may be similarly used.

In a separate embodiment, the enteric coating comprises:
15 (a) cellulose acetate phthalate, (C-A-P) cellulose acetate trimellitate (C-A-T), Hydroxypropyl Methylcellulose Acetate Succinate (HPMCAS), Hydroxypropyl Methylcellulose phthalate (HPMCP), polyvinyl acetate phthalate (PVAP), Anionic phthalate polymers based on methacrylic acid and
20 methacrylic acid esters; (b) compounds either alone or in any combination in organic solvent, (i.e isopropyl alcohol, methanol, ethanol or ethyl acetate) containing at least one plasticizer (i.e. triethyl citrate, polyethylene glycol 6000 or glycerol monostearate, as a coating solution; (c)
25 the group of compounds in (a) either alone or in any combination in aqueous dispersion containing at least one plasticizer can be an alternative coating solution; and (d) the process to apply the coating solution or dispersion to said capsules or tablets is a conventional pharmaceutical
30 method. The coating procedures are performed in a suitable coating machine.

This invention provides a process for manufacturing the capsule dosage form of above described drug which comprises
35 steps of: (a) preparing the granulation by preferably dry mixing the active ingredient and alkaline substance(s) or by non-aqueous wet granulation method using only the

-14-

pharmaceutically acceptable organic solvent, preferably methanol, ethanol or isopropyl alcohol as wetting solution and drying the granulation; (b) filling the capsule with the dried granulation; and (c) coating the capsules with an enteric coating solution or dispersion solution described above.

This invention provides a aqueous-free process for manufacturing the tablet dosage form that comprises steps of (a) preparing the granulation by preferably dry mixing the active ingredient and alkaline substance or pharmaceutical excipients; (b) directly compressing the granulation into a tablet by a conventional method; (c) filling the tablet into an empty hard gelatin capsule shell and; (d) coating the capsule with an enteric coating.

This invention provides an improved pharmaceutical dosage form for oral administration to human being or animal host containing an acid-unstable drug active ingredient which comprises: (a) a core tablet formed by dry mixing drug or its salt with alkaline substance and pharmaceutical excipient or excipients and directly compressing, which can be coated with a protective layer as a barrier to: (i) separate the acid-unstable active drug in the core from the outer enteric coating; and (ii) protect the entered coating from the permeation of alkaline solution formed by water in the core tablet; and (b) an enteric coating disposed on said protectively coated tablet to protect it from releasing the acid-unstable drug in the stomach and to deliver the active to the small intestine. In an embodiment of the above process for manufacturing the improved pharmaceutical dosage form, the core granulation is prepared by dry mixing, direct tablet compressing, subcoating the tablet with organic solvent base protective coating; and coating the subcoated tablets with an enteric coating solution or dispersion.

- 15 -

This invention provides an improved pharmaceutical dosage form for oral administration to human being or animal host containing an acid-unstable drug active ingredient which comprises: (a) a core tablet formed by dry mixing drug or its salt with alkaline substance and pharmaceutical excipient or excipients and directly compressing, which can then be coated with a protective layer as a barrier to: (i) separate the acid-unstable active drug in the core from the outer enteric coating; and (ii) protect the enteric coating from the permeation of alkaline solution formed by water in the core tablet; (b) an enteric coating disposed on said protectively coated tablet to prevent it from releasing the acid-unstable drug in the stomach and to deliver the active to the small intestine; and (c) this tablet is then filled into an empty hard gelatine capsule shell to form a final dosage form. In an embodiment of the above process for manufacturing the improved pharmaceutical dosage form, the core granulation is prepared by dry mixing, direct tablet compressing, subcoating the tablet with organic solvent base protective coating, coating the subcoated tablets with an enteric coating solution or dispersion and filling the said enteric coated tablet into an empty hard gelatin capsule.

Example 1.

Core Granulations: Non-Aqueous Wet Granulation:

- 30 A. Ten grams of Omeprazole were granulated with 10 ml. of ethyl alcohol with agitation. The moist granules were dried and screened to obtain a uniform granule size.
- 35 B. A suspension of Omeprazole being 10 grams in 50 ml. of ethyl alcohol was added into 100 grams of an alkaline inert compound being calcium carbonate granules (DELAVAL, Philadelphia, PA) with agitation to mix homogeneously the liquid and solids. The moist

-16-

granules were dried and screened for uniform and adequate granule size.

- 5 C. Same as Example 1.B., except the alkaline substance is calcium carbonate 90A (Particle Dynamics, Inc., St. Louis, MO.)
- 10 D. A mixture of 16 grams of Omeprazole and 10 grams of povidone USP was dispersed in 15 ml. of ethyl alcohol. The rest of the procedure is same as Example 1.B. above except that the dispersion was used to replace the suspension for 84 grams of alkaline substance.
- 15 E. Ten grams of sodium carboxymethylcellulose was dispersed in 10 ml. of ethyl alcohol. This liquid was then used to granulate a mixture of 2 grams of Omeprazole and 5 grams of calcium carbonate 90A (same as that used in Example 1.C.). The remaining portion of the procedure is the same as Example 1.B. above.
- 20

Example 2.

Core Granulation: Dry Mixing:

- 25 A. Ten grams of Omeprazole were mixed with an alkaline substance being tricalcium phosphate anhydrous USP/NF and then passed through a screen to obtain a homogenous granule size.
- 30 B. Same as Example 2.A. above except the alkaline substance is pharmaceutical excipient microcrystalline cellulose USP/NF.
- 35 C. Same as Example 2.A. above except the alkaline substance is pharmaceutical excipient lactose, anhydrous USP.

-17-

- D. Same as Example 2.A. above except the alkaline substance is pharmaceutical excipient maltodextrin.
- 5 E. Same as Example 2.A. above except the alkaline substance is Calcium Carbonate 90A (same as Example 1.C. above).
- 10 F. Same as Example 2.A. above except the alkaline substance is Calcium Carbonate granules (same as Example 1.B. above).
- G. Same as Example 2.A. above except the alkaline substance is Sodium carboxymethylcellulose.

15

Example 3.

Encapsulation:

- 20 The individual core granulation was mixed with 2% to 5% talc used as a lubricant, and then quantitatively encapsulated in hard gelatin capsules by known pharmaceutical techniques.

25

Example 4.

Direct Tablet Compression:

30

- A. The individual core granulation was mixed with Lactose and Talc or magnesium stearate, and compressed into tablets by known pharmaceutical techniques.